Cosmic Inflation: From Observations to Particle Models

Saturday 18 June 2022 - Saturday 18 June 2022 KEDEA A.U.Th



Book of Abstracts

Contents

Gravitational waves	1
Gravitational Anomalies, torsion and potential geometric origin of the Universe Dark Sector	1
CMB polarization B-mode search with QUBIC and CMB-S4	1
Quintessential Inflation Latest	1
Looking for torsional modified gravity signatures in inflationary observables	1
Modified Gravity Effects on Primordial Gravitational Waves	1
Rescaled Einstein-Hilbert Gravity: Inflation and the Swampland Criteria	1
Cosmic Inflation and Gravity Waves	2
de Sitter vacua, moduli stabilisation and hybrid inflation in string theory	2
Formulating E- & T-Model Inflation in Supergravity	2
New inflationary solutions from old ones	2
Cosmological Hyperfluids, Torsion and Non-metricity	2
Reduced Einstein-Hilbert action: Inflation and the Swampland criteria	2
TBA	3
Models for Freeze-in Barvogenesis	3

1

Gravitational Waves

2

Gravitational Anomalies, torsion and potential geometric origin of the Universe Dark Sector

Corresponding Author: nikolaos.mavromatos@cern.ch

3

CMB polarization B-mode search with QUBIC and CMB-S4

Corresponding Author: s.loucatos@cea.fr

4

Quintessential Inflation Latest

5

Looking for torsional modified gravity signatures in inflationary observables

Corresponding Author: msaridakis@gmail.com

6

Modified Gravity Effects on Primordial Gravitational Waves

8

Rescaled Einstein-Hilbert Gravity: Inflation and the Swampland Criteria

Author: Achilles Gitsis1

Corresponding Author: agitsis62@gmail.com

In this work, a class of $f(R,\phi)$ gravity models is studied which during the inflationary era, which is the large curvature regime, result to an effective inflationary Lagrangian that contains a rescaled Einstein-Hilbert term αR in the presence of a canonical minimally coupled scalar field. The dimensionless parameter α is chosen to take values in the range $0 < \alpha < 1$ and the main motivation for studying these rescaled Einstein-Hilbert $f(R,\phi)$ gravities, is the fact that the rescaled action may render an otherwise incompatible canonical scalar field theory with the Swampland criteria, to be compatible with the Swampland criteria. As it is shown, by studying a large number of inflationary potentials appearing in the 2018 Planck collaboration article for the constraints on inflation, the simultaneous compatibility with both the Planck constraints and the Swampland criteria, is achieved for some models, and the main characteristic of the models for which this is possible, is the small values that the parameter α must take.

9

Cosmic Inflation and Gravity Waves

10

de Sitter vacua, moduli stabilisation and hybrid inflation in string theory

11

Formulating E- & T-Model Inflation in Supergravity

Corresponding Author: kpallis@gen.auth.gr

12

New inflationary solutions from old ones

Corresponding Author: anpaliat@phys.uoa.gr

13

Cosmological Hyperfluids, Torsion and Non-metricity

¹ Aristotle University of Thessaloniki

Reduced Einstein-Hilbert action: Inflation and the Swampland criteria

 $\textbf{Corresponding Author:} \ agits is 62@gmail.com$

15

TBA

 ${\bf Corresponding\ Author:\ kdialekt@gmail.com}$

16

Models for Freeze-in Baryogenesis